arXiv:astro-ph/0410257v1 11 Oct 2004

Looking for outflows from brown dwarfs

E. Masciadri[1]
Max-Plank Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany
`e-mail: masciadr@mpia.de`

A.C. Raga[2]
Ciencias Nucleares, UNAM, Mexico D.F., Mexico

ABSTRACT

First evidences of IR excess and disk mass accretion (strong Hα emission) around brown dwarfs seem to indicate the existence of circumstellar disks around these sub-stellar objects. Nothing is known at the present time about outflows which potentially might be launched from brown dwarfs, although jets are typically associated with the accretion in standard T Tauri star disks. In this paper we calculate the Hα emission of internal working surfaces produced by a radiative jet in a neutral and in a photoionized environment as a function of the jet parameters (the ejection velocity v_j, shock velocity v_s, mass loss rate $\dot{M}$ and radius r_j of the jet) and we provide estimates of the Hα luminosity for the parameters of "standard" Herbig-Haro (HH) jets from T Tauri stars and for the parameters expected for jets from BDs. Interestingly, we find that while the mass loss rates associated with jets from BDs are found to be two orders of magnitude lower than the mass loss rates associated with "standard" HH jets (from T Tauri stars), their velocities are likely to be similar. Based on our calculations, we discuss the conditions in which jets from BDs can be detected and we conclude that the Hα luminosities of internal working surfaces of jets from BDs in a photoionized environment should have only one order of magnitude lower than the Hα luminosities of T Tauri jets in a neutral environment.

Subject headings: ISM: jets and outflows - ISM: Herbig-Haro objects - stars: low-mass, brown dwarfs

1. Introduction

Following the discovery (Nakajima et al. 1995) of the first brown dwarf (BD), several others have been detected in the past few years (Martín et al. 1997; Kirkpatrick et al. 1999, 2001; Gizis et al. 2001; Burgasser et al. 2003; Gizis et al. 2003). Many of the properties of BDs are still unknown, and even the definition of a brown dwarf remains a subject of discussion. One of the fundamental open questions related to BDs is their origin. Are they the result of a molecular cloud collapse followed by the accretion of circumstellar disks formed around a central nucleus (Elmegreen 1999) or are they stellar embryos that, due to dynamical interactions, are ejected from a multiple system before having time to accrete sufficient material from a circumstellar disk to reach the hydrogen burning limit (Reipurth & Clarke 2001; Bate, Bonnel & Bromm 2003) ?

It is even more difficult to define which are the observational evidences that could be used to discern between the two models. At the present time, there are no elements that can exclude one of these models, and it is quite important to collect observational evidence and/or theoretical arguments that can support or reject one or more of these models. It is of course also possible that more than one of the proposed mechanisms might be active in BD formation.

What about the existence of jets from brown dwarfs ? The existence of disks is a necessary condition for the existence of jets. Evidence of accretion disks around BDs was recently obtained by IR excess estimations (IR excesses being the typical signature of the presence of disks around T Tauri stars). Near-IR (3.8 μm) detections in IC348 and Taurus (Liu et al. 2003), mid-IR ($7-14$ μm) detections in ρ Oph and Chamaeleon (Comerón et al. 1998; Comerón et al. 2000; Natta & Testi 2001; Testi et al. 2001; Natta et al. 2002) and sub-millimeter and millimeter detections (André & Montmerle 1994; Carpenter 2002; Klein et al. 2003) started to identify a rich sample of BDs with circumstellar disks. Besides this, signatures of mass outflows are also obtained by the detection of strong Hα emission principally in young BDs (with ages of a few Myr) in the σ Ori region (Barrado y Navascués et al. 2002; Zapatero Osorio et al. 2002), Taurus (Martín et al. 2001), and the TW Hydrae association (Gizis 2002).

It is more difficult to find a straight correlation between the presence of jets and the formation mechanism of BDs. In the scenario in which BDs form from a direct collapse of a molecular cloud, they should have an accretion disk similar to the one around a T Tauri star, and therefore the production of jets would be expected. In the scenario in which BDs are stellar embryos ejected from a young, multiple system, the BDs might still have low mass disks around them, which potentially could also eject bipolar jet systems. Due to the fact that the possible outflow ejection mechanisms associated with these different BD formation models have not yet been studied in any detail, it is not possible to say what a discovery of jets from BDs would imply about the formation mechanism(s) of BDs.

A paper that is particularly relevant in the context of jets from BDs is the one of Quillen & Trilling (1998), who showed that even proto-Jupiters might be expected to produce jets. There is also the work of Wolk & Beck (1990), who extrapolated the outflow luminosity vs. source mass correlation observed for jets from T Tauri (and more massive) stars to low stellar masses, to obtain predictions of the luminosities expected for jets from BDs.

Which is the typical velocity of ejection of BD outflows ? Is the hydrodynamic process producing shock emission in BD jets similar to the one of HH jets from T Tauri stars ? Which is the typical luminosity of their emission ? Can we hope to detect BD jets ? What would be the best way to search for them ?

In the present paper, we discuss a simple, theoretical "internal working surface" model, from which we obtain predictions of the Hα luminosity of an HH jet travelling within a neutral medium or within a photoionized region (section 2). We use this model to obtain predictions of the Hα luminosity of jets from BDs (section 3). Finally, we use our results to propose observational strategies for detecting jets ejected by BDs (section 4).

2. An internal working surface

2.1. The basic geometry

We model the knots along HH jets as "internal working surfaces" resulting from a time-variability of the ejection velocity (Raga et al. (1990), Raga & Kofman (1992), Kofman & Raga (1992)).

In order to obtain analytic estimates of the emission expected from an internal working surface (IWS), we consider the most simple possible analytic model. The schematic diagram of Fig.1 shows the flow structure of an IWS when seen from a frame of reference moving with the velocity v_{ws} of the working surface. In this frame of reference, the material from the upstream and downstream continuous jet segments converges onto the working surface, producing a two-shock structure.

The most simple situation is found for the case in which the upstream and the downstream jet segments have the same density ρ_j. In this case, the material in both beam segments converges onto the working surface with the same velocity v_s (see, e. g., Raga & Kofman 1992).

If the two shocks are strong and radiative, the shock jump conditions can be written as :

$$\rho_j {v_s}^2 = \rho_s {c_s}^2 \,, \tag{1}$$

where ρ_j is the density of the continuous beam segments (see Fig.1), v_s is the shock velocity, and ρ_s and c_s are, respectively, the density and isothermal sound speed (≈ 10 km s^{-1}) at the end of the postshock cooling regions.

Following Falle & Raga (1993), we can now estimate the separation d between the two IWS shocks (see Fig.1) by assuming that the material escapes sideways from the jet beam with a velocity equal to c_s, and then write the inflowing/outflowing mass balance of the IWS :

$$\dot{M}_{in} = \dot{M}_{out}\,, \tag{2}$$

where $\dot{M}_{in} = 2\pi {r_j}^2 \rho_j v_s$ is the rate of mass coming into the IWS through the two shocks, and $\dot{M}_{out} = 2\pi r_j d \rho_s c_s$, where r_j is the jet radius (see Fig.1). From equation (2) we then obtain that the separation d between the two IWS shocks is given by:

$$d = \left(\frac{c_s}{v_s}\right) r_j\,. \tag{3}$$

2.2. Internal working surfaces in a neutral environment

Let us now estimate the Hα luminosity of an IWS of a jet launched in a neutral region. For computing a simple estimate of the emission, we follow Raga & Kofman (1992), and assume that the emission comes mostly from the two shocks within the jet beam, and that the emission from the bow shock (which is pushed into the surrounding environment, see Fig.1) is negligible.

The total Hα luminosity of the IWS is given by :

$$L^{(N)}_{H\alpha} = 2\pi {r_j}^2 \sigma_{H\alpha} \tag{4}$$

where $\sigma_{H\alpha}$ is the luminosity per unit area (of one of the shocks), r_j is the jet radius, and the contribution from the two IWS shocks has been considered.

If one considers the Hα emission from the "self consistent preionization" plane-parallel shock models of Hartigan, Hartmann & Raymond (1987), one finds that for the shock velocity range $v_s = 30 \to 100$ km s^{-1} the Hα luminosity per unit area $\sigma_{H\alpha}$ of the shocks can be fitted with the interpolation formula

$$\sigma_{H\alpha} = 2\times 10^{-5}\,\mathrm{erg\,cm\,s^{-1}}\left(\frac{v_s}{100\,\mathrm{km\,s^{-1}}}\right)^3 n_j\,, \tag{5}$$

where v_s is the shock velocity and n_j ($\approx \rho_j/1.3 m_H$ in our case, see Fig.1) is the preshock number density.

We now write the average mass loss rate from the time-dependent source as

$$\dot{M} = \pi {r_j}^2 \rho_j v_j\,, \tag{6}$$

where v_j is an appropriately defined average value of the ejection velocity. Combining equations (4-5-6) we finally obtain the Hα luminosity

$$L^{(N)}_{H\alpha} = 3.0\times 10^{-3}\mathrm{L}_\odot \left(\frac{\dot{M}}{10^{-7}\mathrm{M}_\odot\mathrm{yr}^{-1}}\right)\times$$

$$\left(\frac{100\,\mathrm{km\,s^{-1}}}{v_j}\right)\left(\frac{v_s}{100\,\mathrm{km\,s^{-1}}}\right)^3\,, \tag{7}$$

normalized to typical parameters for a "normal" HH flow from a T Tauri star.

2.3. Internal working surfaces in a photoionized environment

Let us now estimate the Hα luminosity of an IWS of a jet immersed in a photoionized region. As has been discussed by Raga et al. (2000) and by López-Martín et al. (2001), in many cases HH jets can be completely photoionized by the ionizing radiation field from the H II region source. We will therefore assume that we have an IWS which is fully photoionized by the external, ionizing field.

We will then compute an estimate of the Hα emission coming from the volume of gas in between the two IWS beam shocks (see Fig.1) as :

$$L^{(P)}_{H\alpha} = (\pi r_j^2 d) {n_s}^2 \alpha_{H\alpha}(T_s)\, h\nu_{H\alpha}\,, \tag{8}$$

where the first factor in parentheses on the right is the volume of emitting material (see Fig.1), $n_s = \rho_s/(1.3\,m_H)$ is the post-cooling region number density, $\alpha_{H\alpha}(T_s)$ is the effective Hα recombination coefficient (calculated at the $T_s \approx 10^4$ K temperature of the post-cooling region photoionized material) and $h\nu_{H\alpha}$ is the energy of the Hα transition. Using equations (1), (3) and (6) and setting $c_s \approx 10$ km s^{-1} we then obtain

$$L^{(P)}_{H\alpha} = 1.68\,\mathrm{L}_\odot \left(\frac{\dot{M}}{10^{-7}\mathrm{M}_\odot\mathrm{yr}^{-1}}\right)^2\times$$

$$\left(\frac{100\,\mathrm{km\,s^{-1}}}{v_j}\right)^2\left(\frac{v_s}{100\,\mathrm{km\,s^{-1}}}\right)^3\left(\frac{100\,\mathrm{AU}}{r_j}\right)\,, \tag{9}$$

normalized to typical parameters for a "normal" HH flow from a T Tauri star.

Actually, to this Hα luminosity we should add the contribution from the cooling regions right behind the two IWS shocks. However, from a comparison of equations (9) and (7) we can see that this contribution has to be negligible, at least for the parameters of a "normal" HH jet.

Another interesting feature of fully photoionized jets is that the emission from the jet beam itself can be quite substantial. It is straightforward to show that for a constant density jet the Hα luminosity from the whole of the jet beam is of the order of $L^{(P)}_{H\alpha}$ (the luminosity of the working surface, see equation 9) multiplied by a factor $(L/r_j)(c_s/v_s)^3$ (where L is the total length of the jet beam and $c_s \approx 10$ km s^{-1} is the sound speed of the photoionized gas). For example, for $L/r_j \sim 100$ and $v_s \sim 100$ km s^{-1}, we would have a jet beam Hα luminosity of 10 % of the working surface luminosity. Though being quite diffuse, this emission from the jet beam might also be detectable.

3. Optical emission of outflows from brown dwarfs

In this section we use the results obtained in sections 2.2 and 2.3 to try to compare the emission properties of typical HH jets with the properties of jets ejected by brown dwarfs.

All of the jet (or wind) ejection mechanisms have in common the fact that the ejection velocity v_j is of the order of the escape velocity v_{es} ($v_j \sim v_{es}$) from the surface of the outflow source. For an outflow from a T Tauri star we then have that:

$$v_{es,TT} = \left(\frac{2GM_{TT}}{r_{TT}}\right)^{0.5} , \qquad (10)$$

where M_{TT} is the mass and r_{TT} the radius of a T Tauri star and G is the gravitational constant. For a BD, we will have an outflow with a velocity of the order of the escape velocity

$$v_{es,BD} = v_{es,TT} \cdot \left(\frac{M_{BD} \cdot r_{TT}}{M_{TT} \cdot r_{BD}}\right)^{0.5} , \qquad (11)$$

where M_{BD} is the mass and r_{BD} the radius of a BD.

Which are the typical values for M_{BD} and r_{BD} ? Under the assumption that the BDs which are most likely to produce observable jets are the youngest and most massive ones we can reasonably estimate a range of $M_{BD} \sim 0.03 - 0.075$ M$_\odot$ for the mass. It is more difficult to identify a typical value for r_{BD}. From cool, sub-stellar object atmosphere models (Baraffe et al. 1998; Baraffe et al. 2003; Burrows et al. 1997; Burrows et al. 2001) one sees that for old objects (older than $\sim$ 1 Gyr) the radius of brown dwarfs r_{BD} converges to a typical value of $\sim 0.1\ R_\odot$. For younger BDs (of a few Myr), r_{BD} varies within the $0.1 - 0.4\ R_\odot$ range.

If we then consider typical values $r_{TT} \sim 4\ R_\odot$, $M_{TT} = 1$M$_\odot$ for a T Tauri star, and $M_{BD} = 0.06$M$_\odot$, $r_{BD} \sim 0.3\ R_\odot$ for a BD, from equation (11) we obtain that $v_{es,BD} \sim v_{es,TT}$. Therefore, using typical masses and radii, we find that the escape velocities from the surfaces of BDs have values that are similar to the ones of T Tauri stars. We conclude that the velocities $v_{j,TT}$ and $v_{j,BD}$ of ejection from T Tauri stars and BDs are also comparable.

Another important parameter is of course the shock velocity v_s of the shocks associated with the IWS (see, equations 7 and 9). This shock velocity is not only important for determining the Hα luminosity of the knots, but it also determines the excitation/ionization of the emitted spectrum. In models of outflows from time-dependent sources the shock velocity v_s has a maximum possible value $v_s \sim \Delta v_j/2$ (where Δv_j is the full amplitude of the ejection velocity variability).

As no clear model for the production of ejection variabilities has yet been studied in sufficient detail, it is not possible to find a theoretical relation between the ejection velocity v_j and the amplitude Δv_j of the ejection velocity time-variability. However, from observations of HH jets (see, e. g., the compilation of Raga, Böhm & Cantó 1996) it is clear that the shock velocities of the shocks associated with these objects range from $v_{s,min} \sim 20$ km s^{-1} (this lower limit probably being due to the fact that lower velocity shocks produce very faint line emission and are probably not detected) up to $v_{s,max} \sim 100$ km s^{-1}. Therefore, the amplitude of the velocity variability ranges from low values up to $\Delta v_{j,max} \sim 2\,v_{s,max} \sim 200$ km s^{-1}. In other words, the amplitude of the velocity variability reaches values similar to the mean velocity v_j of the outflow. It appears thus to be reasonable to assume that this result might also be applicable for jets ejected by BDs rather than by T Tauri stars.

Therefore, jets from brown dwarfs have probably knots with emission spectra of similar excitation/ionization to the ones of jets from T Tauri jets. Also, the Hα luminosities of the knots will be given by equations (7) and (9) with values of v_j and v_s similar to the ones given for T Tauri jets.

However, the Hα luminosity is strongly affected by the fact that the accretion rates onto young brown dwarfs appears to be roughly two orders of magnitude lower than the accretion rates onto T Tauri stars (Muzerolle et al. 2003; Barrado y Navascués et al. 2004). One can roughly consider a range of $\dot{M}_{TT} \sim 10^{-8} - 10^{-7}$ M$_\odot$ yr^{-1} for T-Tauri stars and a range of $\dot{M}_{BD} \sim 10^{-10} - 10^{-9}$ M$_\odot$ yr^{-1} for brown dwarfs. As the mass loss rate in an outflow appears to be proportional to the mass accretion rate (Cabrit et al. 1990), this result directly implies that the mass loss rate of a brown dwarf jet is at least two orders of magnitude lower than the mass loss rate of a typical T Tauri jet.

This lower mass loss rate implies that the Hα luminosity of a jet from a BD embedded in a neutral region will be two orders of magnitude lower than the Hα luminosity of a typical jet from a T Tauri star (see equation 7). Taking values of v_j=100 km s^{-1}, v_s=100 $km\ s^{-1}$ and $\dot{M}$=10^{-9} M$_\odot$ yr^{-1} the knots along a brown dwarf jet in a neutral region should have Hα luminosities $L^{(N)}_{H\alpha}(BD) \sim 3 \times 10^{-5}$ L$_\odot$ (see equation 7).

It is not possible to make such a firm statement for the case of a BD jet embedded in a photoionized region. The first thing that we see from equation (9) is that a drop of two orders of magnitude in the mass loss rate results in a drop of four orders of magnitude in the Hα luminosity (when going from a T Tauri to a BD jet). If we assume that the radius of a BD jet has a value similar to the one of a T Tauri jet, we would conclude that the knots along a photoionized BD jet should have an Hα luminosity $L^{(P)}_{H\alpha}(BD) \sim 1.7 \times 10^{-4}$ L$_\odot$ (see equation (9)).

4. Conclusions

We have obtained simple, analytic estimates of the Hα luminosities of internal working surfaces of jets in a neutral environment (see section 2.2) and of fully photoionized jets (see section 2.3).

We have argued that the dynamical characteristics (i. e., jet velocity and amplitude of the ejection velocity variability) are likely to be similar for jets from T Tauri stars and jets from BDs (see section 3). Therefore, the lower observed mass accretion rates (which in its turn determine the mass outflow rates) directly imply that "neutral" brown dwarf jets should have Hα luminosities two orders of magnitude lower than the luminosities of "normal" HH jets from T Tauri stars. This result is in agreement with the Hα luminosities calculated for BD jets by Wolk & Beck (1990), who extended to low source masses the $L_{H\alpha}$ vs. M_{source} correlation found for outflows ejected from T Tauri (and more massive) stars.

We find that the observed correlation can be interpreted in terms of our internal working surface model as a sequence of outflows with the same jet velocity v_j and variability amplitude Δv_j, but with different mass loss rates $\dot{M}_j$.

We find that the knots along photoionized BD jets have Hα luminosities $L^{(P)}_{H\alpha}(BD) \sim 10^{-4}$ L$_\odot$ (see section 4). Even though such luminosities are 4 orders of magnitude lower than the ones of photoionized T Tauri jets (see equation 9), they are only lower by an order of magnitude with respect to the luminosities of "neutral" T Tauri jets (see equation 7).

This result indicates that favourable conditions to carry out searches for BD jets can be found in regions with young, low mass stars embedded within an H II region, such as M 42 (Bally & Reipurth 2001), the Pelican Nebula (Bally & Reipurth 2003) or the IC 1396N H II region (Reipurth et al. 2003). However, as is clear from the published images, in such regions it is difficult to detect jets against the bright and inhomogeneous background of the H II region (which many times has ridges or bright edges which might or might not be jets). This problem could be overcome by searching for photoionized BD jets with a large field Fabry-Pérot interferometer, in which even faint jets should clearly stand out from the H II region emission at high radial velocities.

As a final comment, let us point out that in order to fully photoionize an HH jet, it is necessary to have a relatively strong and nearby source of photoionizing radiation. For producing an Hα luminosity of $\sim 1\,L_\odot$ (as predicted for a jet from a T Tauri star, see equation 9), it is necessary to have $\sim 5 \times 10^{45}$ s^{-1} ionizing photons. In order for a working surface to intercept this number of photons, it is necessary to have something like an O5 star (producing $S_* \approx 5 \times 10^{49}$ s^{-1} ionizing photons) within a distance of $\sim 50\,r_j$ or an O9 star (with $S_* \approx 2 \times 10^{48}$ s^{-1}) within a distance of $\sim 20\, r_j$. The same stars could fully photoionize the much more puny BD jets at distances a factor of ~ 100 times larger than the ones quoted above.

E. Masciadri gratefully acknowledges the hospitality of the Instituto de Ciencias Nucleares, UNAM, Mexico D.F. during this work. The work of A.C. Raga was supported by CONACyT grants 36572-E and 41320 and the DGAPA (UNAM) grant IN 112602. We acknowledge the referees (Bo Reipurth and Noam Soker) for valuable commentaries.

REFERENCES

Andre, P. & Montmerle, T. 1994 ApJ, 420, 837

Bally, J., Reipurth, B. 2003, AJ, 126, 893

Bally, J., Reipurth, B. 2001, ApJ, 546, 299

Baraffe, I., Chabrier, G., Barman, T.M., Allard, F. & Hauschildt, P.H. 2003, A&A, 402, 701

Baraffe, I., Chabier, G., Allard, F. & Hauschildt, P.H. 1998, A&A, 337, 403

Barrado y Navascués, D., Mohanty, S. & Jayawardana R. 2004, ApJ, 604, 248

Barrado y Navascués, D., Zapatero Osorio, M.R., Martín, E.L., Béjar, V.J.S., Rebolo, R. & Mundt, R. 2002, A&A, 393, L85

Bate, M.R., Bonnel, I.A. & Bromm, V. 2003, MNRAS, 339, 577

Burgasser A., Kirkpatrick J., Reid I., Brown M., Miskey C. & Gizis J. 2003, ApJ, 586, 512

Burrows, A., Hubbard, W.B., Lunine, J.I. & Liebert, J. 2001, Rev. of Mod. Phys., 73, 3, 719

Burrows, A., Marley, M., Hubbard, W.B., Lunine, J.I., Guillot, T., Saumon, D., Freedman, R., Sudarsky, D. & Sharp, C. 1997, ApJ, 491, 856

Cabrit, S., Edwards, S., Strom, S.E. & Strom, K.M. 1990, ApJ, 354, 687

Carpenter, J.M. 2002, AJ, 124, 1593

Comerón, F., Neuhaüser, R., Kaas, A.A. 2000, A&A, 359, 269

Comerón, F., Rieke, G.H., Claes, P., Torra, J. & Laureijis, R.J. 1998, A&A, 335, 522

Elmegreen, B.G. 1999, ApJ, 522, 915

Falle, S. A. E. G. & Raga, A. C. 1993, MNRAS, 261, 573

Gizis J.E., Reid I.N., Knapp G.R., Liebert J., Kirkpatrick J. D., Koerner D.W., Burgasser A.J. 2003, AJ, 125, 3202G

Gizis J.E. 2002, ApJ, 575, 484

Gizis, J., Kirkpatrick, J.D., Burgasser, A. et al. 2001, ApJ 551, L163

Hartigan, P., Raymond, J. C. & Hartmann, L. W. 1987, ApJ, 316, 323

Kirkpatrick, J.D., Dahn C.C., Monet, D.D., Reid, I.N., Gizis, J.E., Liebert, J.& Burgasser, A.J. 2001, ApJ, 121, 3235

Kirkpatrick, J.D., Davy, J., Reid, I. N., Liebert, J., Cutri, R.C., Nelson, B., Beichman, C.A., Dahn, C.C., Monet, D.G.,Gizis, J.E. & Skrutskie, M.F. 1999, ApJ, 519, 802

Klein, R., Apai, D., Pascucci, I., Henning, Th. & Waters, L.B.F.M. 2003, ApJ, 593, L57

Kofman, L. & Raga, A. C. 1992, ApJ, 390, 359

Liu, M.C., Najita, J. & Tokunaga, A.T. 2003, ApJ, 585, 372

López-Martín, L., Raga, A. C., López, J. A. & Meaburn, J. 2001, A&A, 371, 1118

Martín E.L., Dougados, C., Magnier, E., Ménard, F., Magazzú A., Cuillandre, J.C. & Delfosse, X. 2001, ApJ, 561, L195

Martín E.L., Basri G., Delfosse X. & Forveille T. 1997, A&A, 327, L29

Muzerolle, J., Hillenbrand, L., Calvet, N., Briceo, C. & Hartmann, L. 2003, ApJ, 592, 266

Nakajima, T., Oppenheimer, B.R., Kulkarni, S.R., Golimowski, D.A., Matthews, K. & Durrance, S.T. 1995, Nature, 378, 463

Natta, A. et al. 2002, A&A, 393, 597

Natta, A. & Testi, L. 2001, A&A, 376, L22

Quillen, A.C. Quillen & Trilling, D.E. 1998, ApJ, 508, 707

Raga, A. C., López-Martín, L., Binette, L., López, J. A., Cantó, J., Arthur, S. J., Mellema, G. & Steffen, W. 2000, MNRAS, 314, 681

Raga, A.C., Böhm K.H. & Cantó, J. 1996, RMxAA, 32, 161

Raga, A. C., Kofman, L. 1992, ApJ, 386, 222

Raga, A. C. et al. 1990, ApJ, 364, 601

Reipurth, B., Armond, T., Raga, A. & Bally, J. 2003, ApJ, 593, L47

Reipurth, B. & Clarke, C. 2001, AJ, 122, 432

Testi, L., Natta, A., Oliva, E., D'Antona, F., Comerón, F., Baffa, C., Comoretto, C. & Gennari, S. 2001, ApJ, 571, L155

Zapatero Osorio, M.R., Béjar, V.J.S., Pavlenko, Ya., Rebolo, R., Allende Prieto, A., Martín, E.L., García López & Mundt, R. 2002, A&A, 384, 937

Wolk, S.J. & Beck, S.C. 1990, PASP, 102, 745



Fig. 1.— Internal working surface (IWS) resulting from a variability of the ejection velocity. The diagram shows the flow as seen from a reference system moving with the IWS. In this reference system, the material in the jet beam converges onto the IWS. If the up- and down-stream jet beam segments have the same density ρ_j, the two IWS shocks have the same shock velocity v_s. The shocked material within the working surface (which has a density ρ_s and sound speed c_s) escapes sonically through the sides and interacts with the surrounding environment, forming a bow shock. The radius of the jet is r_j, and the separation between the two IWS shocks is d.

This 2-column preprint was prepared with the AAS LaTeX macros v5.2.